

Section : QA

Q.1 A donation box can receive only cheques of ₹100, ₹250, and ₹500. On one good day, the donation box was found to contain exactly 100 cheques amounting to a total sum of ₹15250. Then, the maximum possible number of cheques of ₹500 that the donation box may have contained, is

Case Sensitivity: No Answer Type: Equal Possible Answer: 12

Given 30 Answer:

Question Type : SA

Question ID : 48916815395 Status : Answered Q.2 If $c = \frac{16x}{y} + \frac{49y}{x}$ for some non-zero real numbers x and y, then c cannot take the value

Question Type: MCQ

Question ID: 48916814815

Not Attempted and **Marked For Review**

Chosen Option: --

Q.3 If $(3+2\sqrt{2})$ is a root of the equation $ax^2+bx+c=0$, and $(4+2\sqrt{3})$ is a root of the equation $ay^2+my+n=0$, where a,b,c,m and n are integers, then the value of $\left(\frac{b}{m}+\frac{c-2b}{n}\right)$ is

Ans X 1. 0

X 2. 3

3. 4

X 4. 1

Question Type: MCQ

Question ID: 48916814236

Status: Answered

Chosen Option: 3

Suppose the medians BD and CE of a triangle ABC intersect at a point O. If area of **Q.4** triangle ABC is 108 sq. cm., then, the area of the triangle EOD, in sq. cm., is

Case Sensitivity: No Answer Type: Equal Possible Answer: 9

Given 81 Answer:

Question Type: SA

Question ID: 48916815412 Status: Answered

Bob can finish a job in 40 days, if he works alone. Alex is twice as fast as Bob and thrice as fast as Cole in the same job. Suppose Alex and Bob work together on the first day, Bob and Cole work together on the second day, Cole and Alex work together on the third day, and then, they continue the work by repeating this threeday roster, with Alex and Bob working together on the fourth day, and so on. Then, the total number of days Alex would have worked when the job gets finished, is

Case Sensitivity: No Answer Type: Equal Possible Answer: 11

Given 11 Answer:

Question Type: SA

Question ID: 48916815406

Status: Answered

Q.6 A glass contains 500 cc of milk and a cup contains 500 cc of water. From the glass, 150 cc of milk is transferred to the cup and mixed thoroughly. Next, 150 cc of this mixture is transferred from the cup to the glass. Now, the amount of water in the glass and the amount of milk in the cup are in the ratio

Ans

1.1:12.10:3

X 3. 10:13

X 4.3:10

Question Type: MCQ

Question ID : 48916813360 Status : Answered

Chosen Option: 1

Q.7 Consider six distinct natural numbers such that the average of the two smallest numbers is 14, and the average of the two largest numbers is 28. Then, the maximum possible value of the average of these six numbers is

Ans

- 1. 22.5
- X 2. 23
- **X** 3. 23.5
- X 4. 24

Question Type: MCQ

Question ID: 48916813866

Status: Answered

Chosen Option: 1

Q.8 Let r be a real number and $f(x) = \begin{cases} 2x - r & \text{if } x \ge r \\ r & \text{if } x < r \end{cases}$. Then, the equation f(x) = f(f(x)) holds for all real values of x where

Ans

- \checkmark 1. $\chi \leq r$
- $\times 2. \times > r$
- \times 3. $x \ge r$
- \times 4. $x \neq r$

Question Type: MCQ

Question ID: 48916814790

Status : Not Answered

Chosen Option: --

Q.9 Two ships are approaching a port along straight routes at constant speeds. Initially, the two ships and the port formed an equilateral triangle with sides of length 24 km. When the slower ship travelled 8 km, the triangle formed by the new positions of the two ships and the port became right-angled. When the faster ship reaches the port, the distance, in km, between the other ship and the port will be

Ans X 1.4

- 1.4
- 2.6
- **3.** 12
- **X** 4.8

Question Type : MCQ

Question ID: 48916813922

Status : Answered

Chosen Option: 4

Q.10 Nitu has an initial capital of ₹20,000. Out of this, she invests ₹8,000 at 5.5% in bank A, ₹5,000 at 5.6% in bank B and the remaining amount at x% in bank C, each rate being simple interest per annum. Her combined annual interest income from these investments is equal to 5% of the initial capital. If she had invested her entire initial capital in bank C alone, then her annual interest income, in rupees, would have been

Ans X 1. 900

₹ 2.800

× 3. 1000

× 4. 700

Question Type: MCQ

Question ID: 48916813951 Status: Answered

Chosen Option: 2

Q.11

The minimum possible value of $\frac{x^2-6x+10}{3-x}$, for x < 3, is

Ans

 \times 1. -2

√ 2. 2

 $\times 4. -\frac{1}{2}$

Question Type: MCQ

Question ID: 48916812663

Status: Answered

Chosen Option: 2

Q.12 In an examination, the average marks of students in sections A and B are 32 and 60, respectively. The number of students in section A is 10 less than that in section B. If the average marks of all the students across both the sections combined is an integer, then the difference between the maximum and minimum possible number of students in section A is

Case Sensitivity: No Answer Type: Equal Possible Answer: 63

Given 61 Answer:

Question Type: SA

Question ID: 48916815408 Status: Answered

If $\left(\sqrt{\frac{7}{5}}\right)^{3x-y} = \frac{875}{2401}$ and $\left(\frac{4a}{b}\right)^{6x-y} = \left(\frac{2a}{b}\right)^{y-6x}$, for all non-zero real values of a and b, then the value of x+y is

Case Sensitivity: No Answer Type: Equal Possible Answer: 14

Given 6 Answer:

Question Type: SA

Question ID: 48916815390 Status: Answered

Q.14 A group of N people worked on a project. They finished 35% of the project by working 7 hours a day for 10 days. Thereafter, 10 people left the group and the remaining people finished the rest of the project in 14 days by working 10 hours a day. Then the value of N is

Ans X 1. 150

X 2.36

3.140

X 4.23

Question Type: MCQ

Question ID: 48916813818 Status: Answered

Chosen Option: 3

Moody takes 30 seconds to finish riding an escalator if he walks on it at his normal Q.15 speed in the same direction. He takes 20 seconds to finish riding the escalator if he walks at twice his normal speed in the same direction. If Moody decides to stand still on the escalator, then the time, in seconds, needed to finish riding the escalator

Case Sensitivity: No Answer Type: Equal Possible Answer: 60

Given 40 Answer:

Question Type: SA

Question ID: 48916814393 Status: Answered

Q.16 In a triangle ABC, AB = AC = 8 cm. A circle drawn with BC as diameter passes through A. Another circle drawn with center at A passes through B and C. Then the area, in sq. cm, of the overlapping region between the two circles is

Ans \times 1. $16(\pi - 1)$

× 2. 32π

✓ 3. 32(π − 1)

 \times 4. 16 π

Question Type: MCQ

Question ID: 48916814691 Status: Not Answered

Chosen Option: --

2.17	Suppose k is any integer such that the equation $2x^2 + kx + 5 = 0$ has no $x^2 + (k-5)x + 1 = 0$ has two distinct real roots for x . Then, the number	
Ans	× 1. 7	
	→ 2.9	
	× 3. 8	
	× 4. 13	
		Question Type : MCQ
		Question ID : 48916814812
		Status : Answered Chosen Option : 2
).18	The arithmetic mean of all the distinct numbers that can be obtained by rearranging the digits in 1421, including itself, is	
lns	X 1. 2442	
	★ 2. 3333	
	★ 3. 2592	
	✓ 4. 2222	
		Question Type : MCQ
		Question ID : 48916814246
		Status : Not Answered Chosen Option :
Q.19	The lengths of all four sides of a quadrilateral are integer val are of length 1 cm, 2 cm and 4 cm, then the total number of fourth side is	ued. If three of its sides
	are of length 1 cm, 2 cm and 4 cm, then the total number of	ued. If three of its sides
	are of length 1 cm, 2 cm and 4 cm, then the total number of fourth side is 1.5 2.4	ued. If three of its sides
	are of length 1 cm, 2 cm and 4 cm, then the total number of fourth side is 1.5 2.4 3.3	ued. If three of its sides possible lengths of the
	are of length 1 cm, 2 cm and 4 cm, then the total number of fourth side is 1.5 2.4 3.3	ued. If three of its sides
	are of length 1 cm, 2 cm and 4 cm, then the total number of fourth side is 1.5 2.4 3.3	Question Type : MCQ Question ID : 48916813903 Status : Not Answered
	are of length 1 cm, 2 cm and 4 cm, then the total number of fourth side is 1.5 2.4 3.3	Question Type: MCQ Question ID: 48916813903
Ans	are of length 1 cm, 2 cm and 4 cm, then the total number of fourth side is 1.5 2.4 3.3 4.6 Two cars travel from different locations at constant speeds. To starting at the same time, they take 1.5 hours if they travel tows hours if they travel in the same direction. If the speed of the slotten the distance traveled, in km, by the slower car when it me	Question Type : MCQ Question ID : 48916813903 Status : Not Answered Chosen Option :
A.ns	are of length 1 cm, 2 cm and 4 cm, then the total number of fourth side is 1.5 2.4 3.3 4.6 Two cars travel from different locations at constant speeds. To starting at the same time, they take 1.5 hours if they travel tows hours if they travel in the same direction. If the speed of the slo	Question Type : MCQ Question ID : 48916813903 Status : Not Answered Chosen Option :
a.20	are of length 1 cm, 2 cm and 4 cm, then the total number of fourth side is 1.5 2.4 3.3 4.6 Two cars travel from different locations at constant speeds. To starting at the same time, they take 1.5 hours if they travel towards then the distance traveled, in km, by the slower car when it metraveling towards each other, is	Question Type : MCQ Question ID : 48916813903 Status : Not Answered Chosen Option :
λ.ns	are of length 1 cm, 2 cm and 4 cm, then the total number of fourth side is 1.5 2.4 3.3 4.6 Two cars travel from different locations at constant speeds. To starting at the same time, they take 1.5 hours if they travel toware hours if they travel in the same direction. If the speed of the slotten the distance traveled, in km, by the slower car when it metraveling towards each other, is 1.100 2.90	Question Type : MCQ Question ID : 48916813903 Status : Not Answered Chosen Option :
λ.ns	are of length 1 cm, 2 cm and 4 cm, then the total number of fourth side is 1.5 2.4 3.3 4.6 Two cars travel from different locations at constant speeds. To starting at the same time, they take 1.5 hours if they travel toware hours if they travel in the same direction. If the speed of the slotten the distance traveled, in km, by the slower car when it metraveling towards each other, is 1.100 2.90 3.150	Question Type : MCQ Question ID : 48916813903 Status : Not Answered Chosen Option :
Ans	are of length 1 cm, 2 cm and 4 cm, then the total number of fourth side is 1.5 2.4 3.3 4.6 Two cars travel from different locations at constant speeds. To starting at the same time, they take 1.5 hours if they travel toware hours if they travel in the same direction. If the speed of the slotten the distance traveled, in km, by the slower car when it metraveling towards each other, is 1.100 2.90	Question Type : MCQ Question ID : 48916813903 Status : Not Answered Chosen Option :
λ.ns	are of length 1 cm, 2 cm and 4 cm, then the total number of fourth side is 1.5 2.4 3.3 4.6 Two cars travel from different locations at constant speeds. To starting at the same time, they take 1.5 hours if they travel toware hours if they travel in the same direction. If the speed of the slotten the distance traveled, in km, by the slower car when it metraveling towards each other, is 1.100 2.90 3.150	Question Type : MCQ Question ID : 48916813903 Status : Not Answered Chosen Option :
λ.ns	are of length 1 cm, 2 cm and 4 cm, then the total number of fourth side is 1.5 2.4 3.3 4.6 Two cars travel from different locations at constant speeds. To starting at the same time, they take 1.5 hours if they travel toware hours if they travel in the same direction. If the speed of the slotten the distance traveled, in km, by the slower car when it metraveling towards each other, is 1.100 2.90 3.150	Question Type: MCQ Question ID: 48916813903 Status: Not Answered Chosen Option:
λ.ns	are of length 1 cm, 2 cm and 4 cm, then the total number of fourth side is 1.5 2.4 3.3 4.6 Two cars travel from different locations at constant speeds. To starting at the same time, they take 1.5 hours if they travel toware hours if they travel in the same direction. If the speed of the slotten the distance traveled, in km, by the slower car when it metraveling towards each other, is 1.100 2.90 3.150	Question Type : MCQ Question ID : 48916813903 Status : Not Answered Chosen Option : Omeet each other after ards each other, but 10.5 ower car is 60 km/hr, ets the other car while Question Type : MCQ

Q.21 A school has less than 5000 students and if the students are divided equally into teams of either 9 or 10 or 12 or 25 each, exactly 4 are always left out. However, if they are divided into teams of 11 each, no one is left out. The maximum number of teams of 12 each that can be formed out of the students in the school is

Case Sensitivity: No Answer Type: Equal

Possible Answer: 150

Given 20 Answer:

Question Type : SA

Question ID : 48916814385 Status : Answered

Q.22 The average of all 3-digit terms in the arithmetic progression 38, 55, 72, ..., is

Case Sensitivity: No Answer Type: Equal Possible Answer: 548

Given 613 Answer:

Question Type: SA

Question ID : 48916815416 Status : Answered